


MASSACHUSETTS INST OF TECH CAMBRIDGE DEPT OF CHEMISTRY F/G 7/3
THE SYNTHESIS OF 3,3-DIFLUOROALLYL DERIVATIVES OF SILICON BY AN--ETC(U)
AUG 80 D SEYFERTH, K R WURSTHORN, T F LIM N00014-76-C-0837
NR-11 NL

UNCLASSIFIED

$$\frac{1}{\Delta_{\text{eff}}} = \frac{1}{\Delta_{\text{eff}}^{\text{eff}}} + \frac{1}{\Delta_{\text{eff}}^{\text{eff}}}$$

END
DATE
FILMED
9 80
DTIC



LEVEL 4

(12) 1/2

OFFICE OF NAVAL RESEARCH

CONTRACT N00014-76-C-0837 ✓

Task No. NR 053-618

TECHNICAL REPORT NO. 11 ✓

THE SYNTHESIS OF 3,3-DIFLUOROALLYL DERIVATIVES OF SILICON BY AN
ORGANOPHOSPHORUS ROUTE

AD A088354

by

Dietmar Seyferth, Karl R. Wursthorn,
Thomas F. O. Lim and Dennis J. Sepelak

(11) 6 Aug 80
12 25

Prepared for Publication
in the
Journal of Organometallic Chemistry

Massachusetts Institute of Technology
Department of Chemistry, 18-411
Cambridge, Massachusetts 02139

August 6, 1980

DTIC
LECTE
AUG 27 1980
C

Reproduction in whole or in part is permitted for any purpose
of the United States Government.

This document has been approved for public release and sale;
its distribution is unlimited.

DDC FILE COPY

220007

80 8 20 115

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 11	2. GOVT ACCESSION NO. AD-A088	3. RECIPIENT'S CATALOG NUMBER 354
4. TITLE (and Subtitle) "The Synthesis of 3,3-Difluoroallyl Derivatives of Silicon by an Organophosphorus Route"		5. TYPE OF REPORT & PERIOD COVERED Technical Report Interim
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Dietmar Seyferth, Karl R. Wursthorn, Thomas F. O. Lim and Dennis J. Sepelak		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C0837
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Massachusetts Institute of Technology Cambridge, MA 02139		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-618
11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Department of the Navy Arlington, VA 22217		12. REPORT DATE August 6, 1980
		13. NUMBER OF PAGES 18
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approval for Public Release, Distribution Unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES To be published in the Journal of Organometallic Chemistry		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) 3,3-Difluoroallyl derivatives of silicon have been prepared by the reaction of difluorochloromethane with ylides of type $\text{Ph}_3\text{P}=\text{CHCH}_2\text{SiR}_3$ ($\text{SiR}_3 = \text{SiMe}_3, \text{SiMe}_2\text{H}, \text{SiMe}(\text{OSiMe}_3), \text{SiMe}(\text{OSiMe}_3)_2$). Also prepared by this procedure was $\text{Me}_3\text{SiCH}_2\text{C}(\text{CH}_3)=\text{CF}_2$, but $\text{Me}_3\text{SiCH}_2\text{C}(\text{C}_6\text{H}_5)=\text{CF}_2$ could not be prepared in this way, probably because the ylide, $\text{Ph}_3\text{P}=\text{C}(\text{C}_6\text{H}_5)\text{CH}_2\text{SiMe}_3$ was not sufficiently basic to deprotonate HCF_2Cl . The thermal stability of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CF}_2$ was assessed and a few of its reactions examined: addition of CCl_2 and CBr_2 (generated via the appropriate Ph-HgCX_2 (Br) and addition-elimination reactions with phenyl- and n-butyllithium to		

DD FORM 1 JAN 73 1473

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

give $\text{Me}_3\text{SiCH}_2\text{CH}=\text{C}(\text{R})\text{F}$ ($\text{R} = \text{Ph}, \text{n-Bu}$).

Self bar

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DDC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or special
A	

Unclassified

THE SYNTHESIS OF 3,3-DIFLUOROALLYL DERIVATIVES OF SILICON BY AN
ORGANOPHOSPHORUS ROUTE.[#]

Dietmar Seyferth*, Karl R. Wursthorn, Thomas F. O. Lim and
Dennis J. Sepelak

Department of Chemistry
Massachusetts Institute of Technology
Cambridge, Massachusetts 02139 (USA)


[#] Preliminary communication: ref. 1.

SUMMARY


3,3-Difluoroallyl derivatives of silicon have been prepared by the reaction of difluorochloromethane with ylides of type $\text{Ph}_3\text{P}=\text{CHCH}_2\text{SiR}_3$ ($\text{SiR}_3 = \text{SiMe}_3, \text{SiMe}_2\text{H}, \text{SiMe}_2\text{OSiMe}_3, \text{SiMe}(\text{OSiMe}_3)_2$). Also prepared by this procedure was $\text{Me}_3\text{SiCH}_2\text{C}(\text{CH}_3)=\text{CF}_2$, but $\text{Me}_3\text{SiCH}_2\text{C}(\text{C}_6\text{H}_5)=\text{CF}_2$ could not be prepared in this way, probably because the ylide, $\text{Ph}_3\text{P}=\text{C}(\text{C}_6\text{H}_5)\text{CH}_2\text{SiMe}_3$, was not sufficiently basic to deprotonate HCF_2Cl . The thermal stability of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CF}_2$ was assessed and a few of its reactions examined: addition of CCl_2 and CBr_2 (generated via the appropriate PhHgCX_2Br) and addition-elimination reactions with phenyl- and *n*-butyllithium to give $\text{Me}_3\text{SiCH}_2\text{-CH}=\text{C}(\text{R})\text{F}$ ($\text{R} = \text{Ph}, \text{n-Bu}$).

INTRODUCTION


In 1972 Mironov and his coworkers (2) reported the hot tube reaction of vinylmethylchlorosilanes, $\text{CH}_2=\text{CHSiMe}_{3-n}\text{Cl}_n$ ($n = 0-3$), with difluorochloromethane. At temperatures of 485-505°C the yield of the products sought, the respective 2,2-difluorocyclopropylsilanes, I, was optimum. Presumably I were formed by a two-step pro-



cess (eq. 1 and 2). However, compounds of type I showed limited




thermal stability and underwent rearrangement to the respective 3,3-difluoroallylsilane under the reaction conditions (eq. 3). The yields



of the rearrangement products were at a minimum at 485-505°C and increased with increasing temperature, until the tube coked up at 590-610°C and II were the only observed volatile products.

3,3-Difluoroallylsilanes of type II were reported to be stable to 600°C for short periods of time, i.e., they survived passage

through a hot tube at that temperature in a stream of nitrogen. At higher temperature ($\sim 680^{\circ}\text{C}$) these silanes lost CF_2 to give the vinylsilane, $\text{CH}_2=\text{CHSiMe}_{3-n}\text{Cl}_n$, and tetrafluoroethylene. The chemistry of 3,3-difluoroallylsilanes of type II was the subject of a further brief study by this Russian group (3). In the main, reactions were carried out at the reactive Si-Cl functionality in compounds of type II where $n = 1, 2$ and 3: reduction with LiAlH_4 , alkylation with CH_3MgI , methanolysis, hydrolysis and fluorination with antimony trifluoride. In all these reactions the $\text{CH}_2\text{CH}=\text{CF}_2$ substituent survived unchanged. Two C=C addition reactions also were reported (eq. 4, 5).








In view of their apparent high thermal stability and their potentially interesting chemistry at the 3,3-difluoroallyl substituent, these silanes seemed worthy of further investigation. However, it was not clear that the hot tube synthesis was the most practical one possible and, in fact, no useful, detailed synthesis of compounds of type II was provided in the cited papers. In any case, we were interested in the possibility of an alternate, directed synthesis of 3,3-difluoroallylsilanes.

We report here concerning a new route to 3,3-difluoroallyl derivatives of silicon which is based on organophosphorus methodology. Also described are some further reactions of 3,3-difluoroallylsilanes.

RESULTS AND DISCUSSION

3,3-Difluoroallyltrimethylsilane was prepared as shown in eq. 6-10:



Equations 6 (4) and 7 (5) represent well-known chemistry. We have reported all details of the preparation of β -silylalkyltriphenylphosphonium iodides and their conversion to the Wittig reagents in a previous paper (6). Full details also have been provided of the preparation of an analogous tin-substituted phosphonium salt, β -trimethylstannylethyltriphenylphosphonium iodide, and its conversion to the ylide, $\text{Ph}_3\text{P}=\text{CHCH}_2\text{SnMe}_3$ (7) and of the application of the latter to the synthesis of $\text{Me}_3\text{SnCH}_2\text{CH}=\text{CF}_2$ (8). Thus the methodology required for the preparation of 3,3-difluoroallylsilanes as shown in eq. 6-10 was well in hand. In fact, the chemistry of eq. 6-10 was much more easily effected in the case of the silicon compounds than in the case of the preparation of $\text{Me}_3\text{SnCH}_2\text{CH}=\text{CF}_2$. First of all, the required iodomethylsilicon compounds are much more readily prepared

from commercially available chloromethyl-silicon compounds. Secondly, in the case of $\text{Ph}_3\text{PCH}_2\text{CH}_2\text{SnMe}_3^+\text{I}^-$ deprotonation had to be effected with lithium diisopropylamide since organolithium reagents such as phenyllithium attacked at tin as well as at the protons α to the phosphonium center. For successful utilization of the ylide, the amine formed in the deprotonation reaction first had to be removed from the reagent at reduced pressure. In contrast, methyllithium reacted solely as a deprotonating agent in the case of $\text{Ph}_3\text{PCH}_2\text{CH}_2\text{SiMe}_3^+\text{I}^-$. No complications due to competing attack at silicon were encountered.


The reaction shown in eq. 10 is based on chemistry developed by Wheaton and Burton (9) in which a phosphorus ylide first deprotonates HCF_2Cl to give difluorocarbene which then reacts with another mole of the ylide to produce the gem-difluoro-olefin and triphenylphosphine. This reaction proceeded very well with $\text{Ph}_3\text{P}=\text{CHCH}_2\text{SiMe}_3$, giving $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CF}_2$ in 90% yield. At first sight, this reaction has the drawback that two moles of the β -silyl ylide and thus of the β -silylalkylphosphonium salt) are required to produce one mole of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CF}_2$. However, one mole of the ylide is converted to the phosphonium chloride, $\text{Ph}_3\text{PCH}_2\text{CH}_2\text{SiMe}_3^+\text{Cl}^-$, which can be recycled. The triphenylphosphine also can be recovered and used to make more $\text{Ph}_3\text{PCH}_3^+\text{Br}^-$.

3,3-Difluoroallylsilanes with reactive functionality on the silicon atom would be of special interest since incorporation of the 3,3-difluoroallyl group into polysiloxane systems might possibly give silicones with interesting chemical and physical properties. For this reason we also prepared the phosphonium halides $\text{Ph}_3\text{PCH}_2\text{CH}_2\text{SiMe}_2\text{H}^+\text{I}^-$, $\text{Ph}_3\text{PCH}_2\text{CH}_2\text{SiMe}_2\text{OSiMe}_3^+\text{I}^-$ and $\text{Ph}_3\text{PCH}_2\text{CH}_2\text{SiMe}(\text{OSiMe}_3)_2^+\text{I}^-$



(6). These salts could be converted to the respective 3,3-difluoroallylsilicon derivatives, $\text{Me}_2\text{HSiCH}_2\text{CH}=\text{CF}_2$, $\text{Me}_3\text{SiOSiMe}_2\text{CH}_2\text{CH}=\text{CF}_2$ and $(\text{Me}_3\text{SiO})_2\text{SiMeCH}_2\text{CH}=\text{CF}_2$, in good yield. Also prepared by this procedure, using $\text{Ph}_3\text{PCH}(\text{CH}_3)\text{CH}_2\text{SiMe}_3^+\text{I}^-$ (6) as the starting material, was $\text{Me}_3\text{SiCH}_2\text{C}(\text{CH}_3)=\text{CF}_2$. The ylide $\text{Ph}_3\text{P}=\text{C}(\text{C}_6\text{H}_5)\text{CH}_2\text{SiMe}_3$ apparently was not sufficiently basic to deprotonate difluorochloromethane, so $\text{Me}_3\text{SiCH}_2\text{C}(\text{C}_6\text{H}_5)=\text{CF}_2$ could not be prepared by this procedure.

While 3,3-difluoroallylsilicon compounds are stable for short contact times at 600°C (2,3), the thermal stability of such compounds in the liquid phase for extended periods was of greater interest. An NMR tube experiment, using $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CF}_2$ in benzene- d_6 solution, showed this compound to be quite stable. It survived successive heating periods of 15 hr. at 150°C , 6 hr. at 175°C and 21 hr. at 220°C without decomposition and only underwent substantial decomposition after it had been heated at 280°C for 16 hr. In other similar experiments, solutions of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CF}_2$, $\text{Me}_2\text{HSiCH}_2\text{CH}=\text{CF}_2$ and $\text{Me}_3\text{SiOSiMe}_2\text{CH}_2\text{CH}=\text{CF}_2$ in carbon tetrachloride were heated first at 130°C for 16 hr. and then at 200°C for 23 hr. At 130°C , only $\text{Me}_2\text{HSiCH}_2\text{CH}=\text{CF}_2$ showed a small degree of decomposition, probably as a result of reaction of its Si-H function with the solvent. All three compounds decomposed in CCl_4 at 200°C .

Only a few reactions of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CF}_2$ have been investigated thus far. The C=C bond of this silane was found to be reactive toward phenyl(trihalomethyl)mercurial-derived dihalocarbene (eq. 11 and 12):



The addition of organolithium reagents to the C=C bond of polyfluoro-olefins, followed by elimination of lithium fluoride, is a well-known reaction (10). Phenyllithium added to $\text{Me}_3\text{SiCH}_2\text{CH=CF}_2$ (eq. 13), but the yield of the addition-elimination product, obtained as a 1:1 mixture of cis and trans isomers, was only 30%. The results obtained with n-butyllithium were even less satisfactory (eq. 14). Alternate modes of attack by these lithium reagents on 3,3-di-



fluoroallyltrimethylsilane are possible, i.e., difluoroallyl displacement by both RLi, α -deprotonation by n-butyllithium, so the low yields of the desired products are not surprising.

Other C=C bond additions to $\text{Me}_3\text{SiCH}_2\text{CH=CF}_2$ should be possible, but such studies are not within the scope of our present interests. We note that we have also been able to develop a route to the isomeric 1,1-difluoroallylsilanes, $\text{R}_3\text{SiCF}_2\text{CH=CH}_2$, based on the reactions of organochlorosilanes with gem-difluoroallyllithium (8, 11).

EXPERIMENTAL

General Comments.

All reactions were carried out in flame-dried glassware under an atmosphere of dry argon or nitrogen. All solvents were rigorously dried, diethyl ether by distillation from lithium aluminum hydride, tetrahydrofuran from sodium benzophenone ketyl.

Infrared spectra were recorded using a Perkin Elmer 457A infrared spectrophotometer, proton NMR spectra using a Varian Associates T60 spectrometer. Chemical shifts are reported in δ units, ppm downfield from internal tetramethylsilane. Internal standards used were tetramethylsilane, chloroform and dichloromethane. Gas-liquid chromatography (GLC) was used in product analysis, yield determinations by the internal standard method and for the isolation of pure product samples for analysis and spectroscopy.

The preparation of the β -silylalkyltriphenylphosphonium iodides has been described in detail in a previous paper (6).

Preparation of 3,3-Difluoroallyl-silicon Compounds.

(a) 3,3-Difluoroallyltrimethylsilane. A 200 ml, three-necked, round-bottomed flask equipped with a mechanical stirrer, a Dry Ice condenser, a pressure-equalizing addition funnel and a gas inlet tube, was flame-dried, flushed with argon while cooling and charged with 14.71 g (30.0 mmol) of $\text{Ph}_3\text{PCH}_2\text{CH}_2\text{SiMe}_3^+\text{I}^-$ and 50 ml of diethyl ether. The resulting slurry was cooled in an ice bath and then 30 mmol of methyllithium in 16.0 ml of diethyl ether was added, dropwise, with stirring, under argon. The reaction mixture turned red immediately. The resulting solution of $\text{Ph}_3\text{P}=\text{CHCH}_2\text{SiMe}_3$ was stirred at room temperature for 1 hr. It then was cooled to 0°C again and 340 ml (gas volume, about 15 mmol) of difluorochloromethane was con-

* In support of these assignments is previously published work on compounds of type ArCH=CF_2 ($\text{Ar} = \text{C}_6\text{H}_5$, $p\text{-FC}_6\text{H}_4$, $p\text{-O}_2\text{NC}_6\text{H}_4$ and 2-furyl). Their proton NMR spectra showed the following spin-spin coupling constants for the -CH=CF_2 group: J_{FH} (cis): range 3.3-5.0 Hz; J_{FH} (trans): range 25-25.7 Hz (12).

bp 83-85°C.

(b) 3,3-Difluoroallyldimethylsilane. The same procedure was used in the reaction of 10 mmol each of methyllithium and $\text{Ph}_3\text{PCH}_2\text{CH}_2\text{SiMe}_2\text{H}^+\text{I}^-$ in diethyl ether medium. The resulting brownish ylide solution was stirred at room temperature for 1 hr., cooled again to 0°C and treated with about 5 mmol of HCF_2Cl (113 ml gas volume). A solid formed immediately, and after the mixture had been stirred for 2 hr., the supernatant liquid was colorless. Filtration was followed by trap-to-trap distillation (0.05 mm Hg, slight heating) of the filtrate. The colorless distillate was concentrated at reduced pressure and analyzed by GLC (4 ft. SE-30 at room temperature). The product, $\text{Me}_2\text{HSiCH}_2\text{CH}=\text{CF}_2$, was present in 83% yield. Pure samples, n_D^{25} 1.3778, were isolated by GLC. Anal. Found: C, 44.52; H, 7.40. $\text{C}_5\text{H}_{10}\text{F}_2\text{Si}$ calcd.: C, 44.08; H, 7.40. IR ($\text{CCl}_4/\text{CH}_2\text{Cl}_2$), cm^{-1} : 2128s(Si-H), 1742s(C=C), 1328s(=CF₂), 1250s(Me₂Si), 1230s(=CF₂) (principal bands).

In support of these assignments, we note that $\nu(\text{C}=\text{C})$ of $\text{CH}_2=\text{CF}_2$ is found at 1730 cm^{-1} (13), of $\text{Me}_3\text{SnCH}_2\text{CH}=\text{CF}_2$ at 1736 cm^{-1} (8). The =CF₂ group has been reported to have characteristic absorptions at 1340 and 1200 cm^{-1} (14).

¹H NMR ($\text{CCl}_4/\text{CH}_2\text{Cl}_2$): δ 0.08 (d, J 4.0Hz, 6H, Me₂Si), 1.28 (d with some fine splitting, J 8Hz, 2H, SiCH₂), 3.64-4.03 (m, 1H, SiH) and 4.03 (AXY₂ pattern, ²J_{HH} 8Hz, ³J_{FH} (cis) 3Hz, ³J_{FH} (trans) 24Hz, 1H, =CH).

(c) 3,3-Difluoroallylpentamethyldisiloxane. Lithium diisopropylamide was used to deprotonate the phosphonium salt rather than methyllithium since methyllithium is known to cleave the Si-O-Si linkage of hexamethyldisiloxane (15).

To a solution of 0.907 g (8.98 mmol) of diisopropylamine in 25 ml of dry diethyl ether was added dropwise at 0°C, under nitrogen, 8 mmol of *n*-butyllithium in 3.33 ml of hexane. The resulting solution was stirred at room temperature for 1 hr.

A 300 ml, three-necked, round-bottomed flask equipped with a mechanical stirrer, a gas inlet tube and a rubber septum was charged with 4.8 g (8 mmol) of $\text{Ph}_3\text{PCH}_2\text{CH}_2\text{SiMe}_2\text{CSiMe}_3^+\text{I}^-$ and 40 ml of dry Et_2O . To this slurry was added the *i*-Pr₂NLi solution, dropwise with stirring, under nitrogen. The resulting red solution was stirred at room temperature for 1 hr. Subsequently, the mechanical stirrer was replaced with a magnetic stirring assembly and the rubber septum with a ground glass stopper. The gas inlet tube was connected to the vacuum line and all volatiles were removed at 0.5 mm Hg. Forty ml of Et_2O was added to the dry residue to dissolve the ylide and the reaction flask was fitted with a mechanical stirrer and a Dry Ice condenser. About 5 mmol (about 110 ml gas volume) of HCF_2Cl was added at 0°C. The reaction mixture turned brown and a precipitate was formed. The mixture was stirred at room temperature for 3 hr. and worked up as in (a) and (b) above. GLC analysis (4 ft. Se-30 at 100°C) showed that 3.92 mmol (98%) of $\text{Me}_3\text{SiOSiMe}_2\text{CH}_2\text{CH}=\text{CF}_2$ was present. Pure samples were isolated by GLC; n_D^{25} 1.3838. Anal. Found: C, 42.90; H, 8.11. $\text{C}_8\text{H}_{18}\text{OF}_2\text{Si}_2$ calcd.: C, 42.82; H, 8.08. IR (CCl_4), cm^{-1} : 1745 s (C=C), 1329s ($=\text{CF}_2$), 1258s (Me_3Si , Me_2Si), 1232s ($=\text{CF}_2$), 1070s (SiOSi). ^1H NMR ($\text{CCl}_4/\text{CH}_2\text{Cl}_2$): δ 0.02 (s, 9H, Me_3Si), 0.03 (s, 6H, Me_2Si), 1.22 (d of t, $^3\text{J}_{\text{HH}}$ 8Hz, $^4\text{J}_{\text{FH}}$ 1.5Hz, 2H, SiCH_2), 4.03 ppm (12 line pattern, $^2\text{J}_{\text{HH}}$ 8Hz, $^3\text{J}_{\text{FH}}$ (cis) 3Hz, $^3\text{J}_{\text{FH}}$ (trans) 24Hz, 1H, =CH).

(d) 3,3-Difluoroallyl-bis(trimethylsiloxy)methylsilane.

The lithium diisopropylamide route was used in the preparation of the ylide from 6.38 g (10.0 mmol) of $\text{Ph}_3\text{PCH}_2\text{CH}_2\text{SiMe}(\text{OSiMe}_3)_2^+\text{I}^-$ using the procedure described in (c). The ylide was redissolved in diethyl ether and 22 mmol of HCF_2Cl was added at 0°C under nitrogen. The dark red solution gradually faded to light amber and a dark oil separated. The mixture was stirred at room temperature for 10 hr. The ether layer was decanted from the oil. Trap-to-trap distillation of volatiles was followed by concentration of the distillate and examination of the liquid residue by GLC (4 ft. 10% SE-30, 80°C). The desired product, $(\text{Me}_3\text{SiO})_2\text{MeSiCH}_2\text{CH}=\text{CF}_2$, was present in 56% yield (2.80 mmol). Samples of pure product, n_D^{25} 1.3880, were collected by GLC. Anal. Found: C, 40.07; H, 8.05. $\text{C}_{10}\text{H}_{24}\text{O}_2\text{F}_2\text{Si}_3$ calcd.: C, 40.23; H, 8.10. IR (liquid film), cm^{-1} : 1744s ($\text{C}=\text{C}$), 1328m ($=\text{CF}_2$), 1260s (Me_3Si , MeSi), 1233s ($=\text{CF}_2$), 1064vs (SiOSi). ^1H NMR ($\text{CDCl}_3/\text{CHCl}_3$): δ 0.05 (s, 3H, SiCH_3), 0.10 (s, 18H, Me_3Si), 1.14-1.42 (d of t, $^3\text{J}_{\text{HH}}$ 8Hz, $^4\text{J}_{\text{FH}}$ 1.75Hz, 2H, SiCH_2), 3.69-4.50 (12 line pattern, $^2\text{J}_{\text{HH}}$ 8Hz, $^3\text{J}_{\text{FH}}$ (cis) 3Hz, $^3\text{J}_{\text{FH}}$ (trans) 25Hz, 1H, $=\text{CH}$).

(e) 3,3-Difluoro-2-methylallyltrimethylsilane.

Using the apparatus and procedure described in (a), 4.83 g (9.64 mmol) of $\text{Ph}_3\text{PCH}(\text{CH}_3)\text{CH}_2\text{SiMe}_3^+\text{I}^-$ in 25 ml of diethyl ether was treated with 10 mmol of methyllithium in 5.5 ml of ether at 0°C . The red-brown ylide solution was stirred at room temperature for 1 hr. and then 5 mmol (about 120 ml gas volume) was added by the usual procedure. The reaction mixture was stirred at room temperature overnight, filtered, and the filtrate was trap-to-trap distilled

at 0.02 mm Hg. GLC examination of the distillate (6 ft. 10% Apiezon L at 70°C) showed the presence of 3.15 mmol (65%) of the desired product, pure samples of which (n^{25}_D 1.3919) were isolated by GLC. Anal. Found: C, 51.31; H, 8.59. $C_7H_{14}F_2Si$ calcd.: C, 51.18; H, 8.59. IR (liquid film), cm^{-1} : 1753s (C=C). 1H NMR ($CCl_4/CHCl_3$): δ 0.02 (s, 9H, Me_3Si), 1.26 (t, J_{FH} 2.4Hz, 2H, $SiCH_2$), and 1.54 ppm (t, J_{FH} 3.2Hz, 3H, $=CCH_3$).

The salts in the filter cake were dried to give 2.47 g of $Ph_3PCH(CH_3)CH_2SiMe_3^+Cl^-$. The residue from the trap-to-trap distillation was recrystallized from ethanol to give 0.91 g (3.47 mmol, 72%) of Ph_3P .

Dihalocarbene Addition to 3,3-Difluoroallyltrimethylsilane.


(a) Dichlorocarbene. A 100 ml one-necked, round-bottomed flask equipped with a reflux condenser topped with a nitrogen inlet tube and a magnetic stir-bar was charged with 2.408 g (5.6 mmol) of $PhHgCCl_2Br$ (16), 0.874 g (5.83 mmol) of $Me_3SiCH_2CH=CF_2$ and 20 ml of dry benzene. The resulting solution was stirred and heated at reflux (ca. 80°C) under nitrogen for 90 min. The reaction mixture was allowed to cool to room temperature and was filtered to remove phenylmercuric bromide. The filtrate was trap-to-trap distilled at 0.05 mm Hg with slight heating into a receiver cooled to -78°C. GLC analysis (4 ft. SE-30 at 110°C) of the distillate showed the presence of 4.31 mmol (77%) of 1,1-difluoro-2,2-dichloro-3-(trimethylsilylmethyl)cyclopropane, n^{25}_D 1.4297. Anal. Found: C, 36.24; H, 5.14. $C_7H_{12}F_2Cl_2Si$ calcd.: C, 36.06; H, 5.19. 1H NMR (CCl_4/CH_2Cl_2): δ 0.13 (s, 9H, Me_3Si), 0.73-1.02 (m, 2H, $SiCH_2$) and 1.67-2.32 ppm (m, 1H, cyclopropyl H).

(b) Dibromocarbene. The same procedure was used in the reaction of 1.827 g (3.45 mmol) of PhHgCBr_3 (16), 0.494 g (3.29 mmol) of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CF}_2$ in 5 ml of benzene at 80°C for 5 hr. GLC analysis of the final trap-to-trap distillate (4 ft. SE-30 at 130°C) showed the presence of 2.43 mmol (70%) of 1,1-difluoro-2,2-dibromo-3-(trimethylsilylmethyl)cyclopropane, n_D^{25} 1.4698. Anal. Found: C, 26.14; H, 3.82. $\text{C}_7\text{H}_{12}\text{F}_2\text{Br}_2\text{Si}$: calcd.: C, 26.10; H, 3.76. ^1H NMR ($\text{CCl}_4/\text{CH}_2\text{Cl}_2$): δ 0.03 (s, 9H, Me_3Si), 0.56-0.89 (m, 2H, SiCH_2) and 1.63-2.31 (m, 1H, cyclopropyl H).

Reaction of 3,3-Difluoroallyltrimethylsilane with Organolithium Reagents.


To a solution of 3.82 mmol of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CF}_2$ in 5 ml of dry diethyl ether (50 ml three-necked flask equipped with magnetic stir-bar, nitrogen inlet tube and addition funnel) was added, dropwise with stirring, 3.82 mmol of freshly prepared phenyllithium in 6.7 ml of 4:1 Et_2O /benzene. The reaction mixture was stirred at room temperature for 17 hr. and the resulting cloudy solution then was trap-to-trap distilled at 0.05 mm Hg (with heating) into a receiver cooled to -78°C . GLC examination of the distillate (6 ft. Apiezon L at 140°C) showed the presence of 1.14 mmol (30%) of a 1:1 mixture of the cis and trans isomers of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CFPh}$. The isomers were separable by GLC and were individually characterized.


$\text{Me}_3\text{SiCH}_2\text{CH}=\text{CFPh}$ n_D^{25} 1.5138. Anal. Found: C, 69.33; H, 8.23. $\text{C}_{12}\text{H}_{17}\text{FSi}$ calcd.: C, 69.17; H, 8.23. ^1H NMR ($\text{CCl}_4/\text{CH}_2\text{Cl}_2$): δ 0.06 (s, 9H, Me_3Si), 1.62 (d of d, J_{HH} 8.5Hz, J_{FH} 1.5Hz, 2H, SiCH_2) and 5.35 ppm (d of t, J_{FH} 36.3Hz, 1H, =CH).



The isomer assignments are based on the the magnitude of J_{FH} in the vinylic system. In published examples, J_{FH} (trans) is larger than J_{FH} (cis) in $\text{RCH}=\text{CHF}$ compounds. For instance, in the cis and trans isomers of $\text{PhCH}=\text{CHF}$ and $n\text{-C}_6\text{H}_{13}\text{CH}=\text{CHF}$, Burton and Greenlimb reported J_{FH} (trans) in the range of 40-44Hz, J_{FH} (cis) in the range 17.9-19.6Hz (17). The geminal coupling constants were in the range 79.5-85.4 $\frac{\text{Hz}}{\text{Hz}}$, thus excluding the alternate structure, $\text{Me}_3\text{SiCH}_2(\text{Ph})\text{C}=\text{CHF}$, for our product.

A similar reaction was carried out with n-butyllithium. The lithium reagent, 4.41 mmol in hexane, was added to 4.41 mmol of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CF}_2$ in 5 ml of diethyl ether at -78°C and then 0.44 mmol of N,N,N',N'-tetramethylethylenediamine was added by syringe. The mixture was allowed to warm to room temperature and then was stirred for 2 hr. Trap-to-trap distillation of volatiles at 0.05 mm Hg with heating was followed by GLC analysis of the distillate (6 ft. Apiezon L at 120°C , 6 ft. silver nitrate at 58°C). The two isomers of $\text{Me}_3\text{SiCH}_2\text{CH}=\text{CF}(\text{n-Bu})$ were present in 20% combined yield. Pure samples of each were isolated by GLC.





(3 parts)

^1H NMR ($\text{CCl}_4/\text{CHCl}_3$): δ -0.12 (s, 9H, Me_3Si), 0.58-1.50 (m, 9H, C_4H_9), 1.08 (d, J_{HH} 9Hz, 2H, SiCH_2) and 4.88 ppm (d of t, J_{FH} 22Hz, 1H, =CH).

The isomer mixture had n_D^{25} 1.4309.

Acknowledgment. This work was supported in part by the Office of Naval Research.

REFERENCES

1. D. Seyferth, K. R. Wursthorn and R. E. Mammarella, J. Org. Chem., 42 (1977) 3104.
2. V. F. Mironov, O. M. Rad'kova, V. D. Sheludyakov and V. V. Shcherbinin, Dokl. Akad. Nauk SSSR, 207 (1972) 114.
3. V. D. Sheludyakov, V. V. Shcherbinin, N. A. Viktorov and V. F. Mironov, Zh. Obshch. Khim., 44 (1974) 1935.
4. F. C. Whitmore and L. H. Sommer, J. Amer. Chem. Soc., 68 (1946) 481.
5. H.-J. Bestmann, Angew. Chem. Int. Ed. Eng., 4 (1965) 583.
6. D. Seyferth, K. R. Wursthorn, T. F. O. Lim and D. J. Sepelak, J. Organometal. Chem., 181 (1979) 293.
7. D. Seyferth, K. R. Wursthorn and R. E. Mammarella, J. Organometal. Chem., 179 (1979) 25.
8. D. Seyferth and K. R. Wursthorn, J. Organometal. Chem., 182 (1979) 455.
9. G. A. Wheaton and D. J. Burton, Tetrahedron Lett. (1976) 895.
10. W. A. Sheppard and C. M. Sharts, "Organic Fluorine Chemistry", Benjamin, New York, 1969, Chapters 3 and 6.
11. D. Seyferth, R. M. Simon, D. J. Sepelak and H. A. Klein, J. Org. Chem., 45 (1980) 2273.
12. S. A. Fuqua, W. G. Duncan and R. M. Silverstein, J. Org. Chem., 30 (1965) 1027.
13. P. Torkington and H. W. Thompson, Trans. Faraday Soc., 41 (1945) 236.
14. R. B. Barnes, R. C. Gore, R. W. Stafford and V. Z. Williams, Analyt. Chem., 20 (1948) 402.

15. D. Seyferth and D. L. Alleston, *Inorg. Chem.*, 2 (1963) 418.
16. D. Seyferth and R. L. Lambert, Jr., *J. Organometal. Chem.*,
16 (1969) 21.
17. D. J. Burton and P. E. Greenlimb, *J. Org. Chem.*, 40 (1975)
2796.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	2	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709	1
ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	1	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152	1
ONR Branch Office Attn: Scientific Dept. 715 Broadway New York, New York 10003	1	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555	1
ONR Branch Office 1030 East Green Street Pasadena, California 91106	1	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1
ONR Branch Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	1	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940	1
Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	1	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380	1
The Assistant Secretary of the Navy (R,E&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	1	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	1	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401	1
Defense Documentation Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232	1
Dr. Fred Saalfeld Chemistry Division Naval Research Laboratory Washington, D.C. 20375	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1

TECHNICAL REPORT DISTRIBUTION LIST, GENNo.
Copies

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco 96503

1

Mr. James Kelley
DTNSRDC Code 2803
Annapolis, Maryland 21402

1

TECHNICAL REPORT DISTRIBUTION LIST, 053

	<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
Dr. R. N. Grimes University of Virginia Department of Chemistry Charlottesville, Virginia 22901	1	Dr. M. H. Chisholm Department of Chemistry Indiana University Bloomington, Indiana 47401	1
Dr. M. Tsutsui Texas A&M University Department of Chemistry College Station, Texas 77843	1	Dr. B. Foxman Brandeis University Department of Chemistry Waltham, Massachusetts 02154	1
Dr. M. F. Hawthorne University of California Department of Chemistry Los Angeles, California 90024	1	Dr. T. Marks Northwestern University Department of Chemistry Evanston, Illinois 60201	1
Dr. D. B. Brown University of Vermont Department of Chemistry Burlington, Vermont 05401	1	Dr. G. Geoffrey Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802	1
Dr. W. B. Fox Naval Research Laboratory Chemistry Division Code 6130 Washington, D.C. 20375	1	Dr. J. Zuckerman University of Oklahoma Department of Chemistry Norman, Oklahoma 73019	1
Dr. J. Adcock University of Tennessee Department of Chemistry Knoxville, Tennessee 37916	1	Professor O. T. Beachley Department of Chemistry State University of New York Buffalo, New York 14214	1
Dr. A. Cowley University of Texas Department of Chemistry Austin, Texas 78712	1	Professor P. S. Skell Department of Chemistry The Pennsylvania State University University Park, Pennsylvania 16802	1
Dr. W. Hatfield University of North Carolina Department of Chemistry Chapel Hill, North Carolina 27514	1	Professor K. M. Nicholas Department of Chemistry Boston College Chestnut Hill, Massachusetts 02167	1
		Professor R. Neilson Department of Chemistry Texas Christian University Fort Worth, Texas 76129	1
Professor H. Abrahamson University of Oklahoma Department of Chemistry Norman, Oklahoma 73019	1	Professor M. Newcomb Texas A&M University Department of Chemistry College Station, Texas 77843	1